

Heavy flavor kinetics at the hadronization transition

D. Blaschke^{a,b}, G. Burau^a, Yu. Kalinovsky^c, T. Barnes^{d,e}

^a Fachbereich Physik, Universität Rostock, Germany

^b Bogolyubov Laboratory for Theoretical Physics, JINR Dubna, Russia

^c Laboratory for Information Technologies, JINR Dubna, Russia

^d Oak Ridge National Laboratory, Oak Ridge (TN), USA

^e Department of Physics, University of Tennessee, Knoxville TN, USA

E-mail: david.blaschke@physik.uni-rostock.de

Abstract. We investigate the in-medium modification of the charmonium breakup processes due to the Mott effect for light (π , ρ) and open-charm (D , D^*) quark-antiquark bound states at the chiral/deconfinement phase transition. The Mott effect for the D -mesons effectively reduces the threshold for charmonium breakup cross sections, which is suggested as an explanation of the anomalous J/ψ suppression phenomenon in the NA50 experiment. Further implications of finite-temperature mesonic correlations for the hadronization of heavy flavors in heavy-ion collisions are discussed.

The J/ψ meson plays a key role in the experimental search for the quark-gluon plasma (QGP) in heavy-ion collision experiments [1] where an anomalous suppression of its production cross section relative to the Drell-Yan continuum as a function of the centrality of the collision has been found by the CERN-NA50 collaboration [2]. An effect like this has been predicted to signal QGP formation [3] as a consequence of the screening of color charges in a plasma in close analogy to the Mott effect (metal-insulator transition) in dense electronic systems [4]. However, a necessary condition to explain J/ψ suppression in the static screening model is that a sufficiently large fraction of $c\bar{c}$ pairs after their creation have to traverse regions of QGP where the temperature (resp. parton density) has to exceed the Mott temperature $T_{J/\psi}^{\text{Mott}} \sim 1.2 - 1.3 T_c$ [5, 6] for a sufficiently long time interval $\tau > \tau_f$, where $T_c \sim 170$ MeV is the critical phase transition temperature and $\tau_f \sim 0.3$ fm/c is the J/ψ formation time. Within an alternative scenario [7], J/ψ suppression does not require temperatures well above the deconfinement one but can occur already at T_c due to impact collisions by quarks from the thermal medium. An important ingredient for this scenario is the lowering of the reaction threshold for string-flip processes which lead to open-charm meson formation and thus to J/ψ suppression. This process has an analogue in the hadronic world, where e.g. $J/\psi + \pi \rightarrow D^* + \bar{D} + h.c.$ could occur provided the reaction threshold of $\Delta E \sim 640$ MeV can be overcome by pion impact. It has been shown recently [8, 9] that this process and its in-medium modification can play a key role in the understanding of anomalous J/ψ suppression as a deconfinement signal. Since at the deconfinement transition the D -mesons enter the continuum of unbound (but strongly correlated) quark-antiquark states (Mott-effect), the relevant threshold for charmonium breakup is lowered and the reaction rate for the process gets critically enhanced. Thus a process which is negligible in the vacuum may give rise to additional (anomalous) J/ψ suppression when conditions of the chiral/deconfinement

transition and D - meson Mott effect are reached in a heavy-ion collision but the dissociation of the J/ψ itself still needs impact to overcome the threshold which is still present but dramatically reduced.

For this alternative scenario as outlined in [9] to work the J/ψ breakup cross section by pion impact is required and its dependence on the masses of the final state D - mesons has to be calculated. Both, nonrelativistic potential models [10, 11] and chiral Lagrangian models [12, 13, 14] have been employed to determine the cross section in the vacuum. The results of the latter models appear to be strongly dependent on the choice of formfactors for the meson-meson vertices. This is considered as a basic flaw of these approaches which could only be overcome when a more fundamental approach, e.g. from a quark model, can determine these input quantities of the chiral Lagrangian approach. Work in this direction is in progress, see [15], but meanwhile we have to rely on phenomenological approaches. Recently, a modified scheme for




Figure 1. Energy dependent breakup cross section for the subprocess $J/\psi + \pi \rightarrow D + \bar{D}^*$ in the nonrelativistic quark exchange model [11] (symbols with error bars) and in the chiral Lagrangian model [16] (solid and dashed lines).

defining formfactors of meson-meson vertices has been suggested [16] which uses Gaussian functions instead of monopole ones [13, 14] and respects differences in the effective ranges of contact and meson exchange diagrams which arise from their quark substructure. The result is shown as the solid line in Fig. 1 and compares well with that of the quark exchange model ([11], filled circles) in the peak value and position. The fast drop of the cross section above the threshold due to the behavior of the meson wave functions is described more appropriately by Gaussian formfactors than by monopole ones, for details see [16]. A change in the coupling constant of the contact term (which corresponds to the capture diagrams in the quark exchange approach) by a factor three leads only to a change in the total cross section by about 50 % (dashed line in Fig. 1). Thus the dominant contributions come from the meson exchange

diagrams, which are related to the transfer diagrams of Ref. [11]. Chiral Lagrangian and quark exchange approaches to the hadronic interactions of charmonium can be brought to good agreement with each other.

Since at the hadronization transition the J/ψ is expected to be a true bound state with almost unchanged mass the most dramatic in-medium modification for J/ψ breakup processes is expected due to the Mott-effect for pions, rho- and D-mesons, i.e. their transition from bound states to resonant scattering states in the quark-antiquark continuum [8, 9]. Formally, this transition can be described by the behavior of the spectral function in corresponding channel. As a result, the kinematics for processes involving heavy mesons is not constrained to their (vacuum) mass shell so that they become *subthreshold*. The result of a model calculation which uses an off-shell extrapolation of the quark exchange cross section [11] is given in Fig. 2. The




Figure 2. In-medium modification of the J/ψ breakup cross section by pion impact due to the Mott effect for open charm mesons at $T = 150$ MeV.

corresponding reaction rate for J/ψ dissociation shows a step function behavior at the Mott (hadronization) transition [9] which has earlier been postulated [17] without a microscopic foundation for a successful description of the anomalous J/ψ suppression pattern. The present approach is a relativistic formulation of string-flip processes which have been suggested for a unified description of J/ψ suppression in hadronic as well as in quark matter [7]. It predicts a simultaneous increase in the reaction rates for breakup of all hidden charm (bottom) states once the Mott effect for the final state open charm (bottom) hadrons occurs. Upon neglect of this trigger mechanism one would predict instead a sequential breakup of heavy quarkonia according to their different dissociation energies into the quark-antiquark continuum [18, 19, 20].

Applications to the description of charmonium production in nucleus-nucleus collisions are in progress [21].

In summary, we have shown that recent developments of the chiral Lagrangian and quark exchange model approaches have lead to a convergence of results for quarkonium-hadron breakup (string-flip) cross sections. Reaction rates for these processes are negligibly small in the hadronic world but show a critical enhancement at the deconfinement/ hadronization transition when the Mott effect for open flavor hadrons is taken into account. This reduces dramatically those reaction thresholds which would otherwise prevent heavy flavor equilibration in hadronic matter. The inclusion of the Mott effect into the description of heavy flavor kinetics at the hadronization transition allows to link three recently observed phenomena: (i) anomalous J/ψ suppression [2, 8, 21], (ii) open charm enhancement [22, 23], and (iii) the success of the statistical model [24, 25, 26].

Acknowledgments

The research of Yu.K. has been supported by DFG grant No. 436 RUS 17/62/01. D.B. and G.B. acknowledge support by DAAD for their visits at Oak Ridge National Laboratory and University of Tennessee at Knoxville. T.B. has been supported in part by NSF grant No. INT-0004089.

References

- [1] Abreu M C *et al.* (NA50 Collaboration) 2000 *Phys. Lett. B* **477** 28
- [2] Abreu M C *et al.* (NA50 Collaboration) 1997 *Phys. Lett. B* **410** 337
Abreu M C *et al.* (NA50 Collaboration) 1999 *Phys. Lett. B* **450** 456
- [3] Matsui T and Satz H 1986 *Phys. Lett. B* **178** 416
- [4] Redmer R 1997 *Phys. Rep.* **282** 35
- [5] Karsch F, Mehr M T and Satz H 1988 *Z. Phys. C* **37** 617
- [6] Röpke G, Blaschke D and Schulz H 1988 *Phys. Lett. B* **202** 479
- [7] Blaschke D, Röpke G and Schulz H 1989 *Phys. Lett. B* **233** 434.
- [8] Blaschke D, Burau G and Kalinovsky Y L 2000 in *Progress in Heavy Quark Physics V*, Ivanov M A, Lyubovitskij V E and Lipartia E (Eds), Dubna, p 144 [arXiv:nucl-th/0006071]
- [9] Burau G R G, Blaschke D B and Kalinovsky Y L 2001 *Phys. Lett. B* **506** 297 [arXiv:nucl-th/0012030]
- [10] Martins K, Blaschke D and Quack E 1995 *Phys. Rev. C* **51** 2723
- [11] Wong C Y, Swanson E S and Barnes T 2000 *Phys. Rev. C* **62** 045201 [arXiv:hep-ph/9912431];
Wong C Y, Swanson E S and Barnes T 2002 *Phys. Rev. C* **65** 014903 [arXiv:nucl-th/0106067].
- [12] Matinyan S G and Müller B 1998 *Phys. Rev. C* **58** 2994
- [13] Lin Z and Ko C M 2000 *Phys. Rev. C* **62** 034903.
- [14] Haglin K L and Gale C 2001 *Phys. Rev. C* **63** 065201.
- [15] Blaschke D B, Burau G R G, Ivanov M A, Kalinovsky Y L and Tandy P C 2000 in *Progress in Nonequilibrium Green Functions*, Bonitz M (Ed), World Scientific, Singapore, p 392 [arXiv:hep-ph/0002047]
- [16] Ivanov V V, Kalinovsky Y L, Blaschke D B and Burau G R G 2001 in *Exploring quark matter*, Burau G, Blaschke D and Schmidt S (Eds), Rostock University Press [arXiv:nucl-th/0111014];
Preprint ECT*-01-34, arXiv:hep-ph/0112354
- [17] Blaizot J P and Ollitrault J Y 1996 *Phys. Rev. Lett.* **77** 1703 [arXiv:hep-ph/9606289];
Blaizot J P, Dinh M and Ollitrault J Y 2000 *Phys. Rev. Lett.* **85** 4012 [arXiv:nucl-th/0007020].
- [18] Nardi M and Satz H 1998 *Phys. Lett. B* **442** 14 [arXiv:hep-ph/9805247].
- [19] Digal S, Petreczky P and Satz H 2001 *Phys. Rev. D* **64** 094015 [arXiv:hep-ph/0106017].
- [20] Wong C Y, Barnes T, Swanson E S and Crater H W 2001 *Preprint* arXiv:nucl-th/0112023
- [21] Blaschke D, Burau G and Polleri A 2001 work in progress.
- [22] Abreu M C *et al.* (NA38 and NA50 Collaborations) 2000 *Eur. Phys. J. C* **14** 443
- [23] Kostyuk A P, Gorenstein M I, Stöcker H and Greiner W 2001 *Phys. Lett. B* **519** 207
- [24] Braun-Munzinger P and Stachel J 2000 *Phys. Lett. B* **490** 196
- [25] Kostyuk A P, Gorenstein M I, Stöcker H and Greiner W 2001 *Preprint* arXiv:hep-ph/0110269
- [26] Grandchamp L and Rapp R 2001 *Preprint* arXiv:hep-ph/0103124.